COUNTY OF APPOMATTOX

Department of Community and Economic Development P. O. Box 787 Appomattox, VA 24522

INSPECTION GUIDELINES


FOR

EROSION & SEDIMENT CONTROL




LONNY HAMLETT, BUILDING OFFICIAL

PHONE: (434) 352-8183 • FAX: (434) 352-4214



Source: Adapted from 1983 Maryland Standards for Soil Erosion and Sediment Control, and Va. DSWC

Plate 3.02-1



Source: Adapted from <u>Installation of Straw and Fabric Filter</u>
Barriers for Sediment Control, Sherwood and Wyant

Plate 3.05-2

TABLE 3.32-D SITE SPECIFIC SEEDING MIXTURES FOR PIEDMONT AREA

| | Total Lbs. Per Acre | | |
|--|---|--|--|
| Minimum Care Lawn | | | |
| Commercial or Residential Kentucky 31 or Turf-Type Tall Fescue Improved Perennial Ryegrass Kentucky Bluegrass | 175-200 lbs. 95-100% 0-5% 0-5% | | |
| High-Maintenance Lawn | 200-250 lbs. | | |
| - Kentucky 31 or Turf-Type Tall Fescue | 100% | | |
| General Slope (3:1 or less) | | | |
| - Kentucky 31 Fescue - Red Top Grass - Seasonal Nurse Crop * | 128 lbs. 2 lbs. 20 lbs. 150 lbs. | | |
| Low-Maintenance Slope (Steeper than 3:1) | | | |
| Kentucky 31 Fescue Red Top Grass Seasonal Nurse Crop * Crownvetch ** | 108 lbs. 2 lbs. 20 lbs. 20 lbs. 150 lbs. | | |
| * Use seasonal nurse crop in accordance with seeding dates as stated below: February 16th through April | | | |
| ** Substitute Sericea lespedeza for Crownvetch east of Farmville, Va. (May through September use hulled Sericea, all other periods, use unhulled Sericea). If Flatpea is used in lieu of Crownvetch, increase rate to 30 lbs./acre. All legume seed must be properly inoculated. Weeping Lovegrass may be added to any slope or low-maintenance mix during warmer seeding periods; add 10-20 lbs./acre in mixes. | | | |

Mulching

- 1. Seedings <u>made in fall for winter cover and during hot and dry summer months</u> shall be mulched according to MULCHING, Std. & Spec. 3.35, except that hydromulches (fiber mulch) will not be considered adequate. Straw mulch should be used during these periods.
- 2. Temporary seedings made under favorable soil and site conditions during optimum spring and fall seeding dates may not require mulch.

Re-seeding

Areas which fail to establish vegetative cover adequate to prevent rill erosion will be reseeded as soon as such areas are identified.

| TABLE 3.31-B ACCEPTABLE TEMPORARY SEEDING PLANT MATERIALS "QUICK REFERENCE FOR ALL REGIONS" | | | |
|---|---|------------------|--|
| Planting Dates | Species | Rate (lbs./acre) | |
| Sept. 1 - Feb. 15 | 50/50 Mix of Annual Ryegrass (Lolium multi-florum) & Cereal (Winter) Rye (Secale cereale) | 50 - 100 | |
| Feb. 16 - Apr. 30 | Annual Ryegrass (Lolium multi-florum) | 60 - 100 | |
| May 1 - Aug 31 | German Millet (Setaria italica) | 50 | |

Source: Va. DSWC

TABLE 3.35-A
ORGANIC MULCH MATERIALS AND APPLICATION RATES

| | RATES: | | | |
|--------------------------------------|---|------------------|--|--|
| MULCHES: | Per Acre | Per 1000 sq. ft. | NOTES: | |
| Straw or Hay | 1½ - 2 tons (Minimum 2 tons for winter cover) | 70 - 90 lbs. | Free from weeds and coarse matter. Must be anchored. Spread with mulch blower or by hand. | |
| Fiber Mulch | Minimum 1500 lbs. | 35 lbs. | Do not use as mulch for winter cover or during hot, dry periods.* Apply as slurry. | |
| Corn Stalks | 4 - 6 tons | 185 - 275 lbs. | Cut or shredded in 4-6" lengths. Air-dried. Do not use in fine turf areas. Apply with mulch blower or by hand. | |
| Wood Chips | 4 - 6 tons | 185 - 275 lbs. | Free of coarse matter. Airdried. Treat with 12 lbs nitrogen per ton. Do not use in fine turf areas. Apply with mulch blower, chip handler, or by hand. | |
| Bark Chips or Shredded Bark | 50 - 70 cu. yds. | 1-2 cu. yds. | Free of coarse matter. Airdried. Do not use in fine turf areas. Apply with mulch blower, chip handler, or by hand. | |

^{*} When fiber mulch is the only available mulch during periods when straw should be used, apply at a minimum rate of 2000 lbs./ac. or 45 lbs./1000 sq. ft.

Source: Va. DSWC